arXiv:1208.2244v1 [math.GM] 9 Aug 2012

Some considerations in favor of the truth of Goldbach's Conjecture

Ciro D'Urso

August 13, 2012

Italian Senate

ciro.durso@senato.it

(paper submitted to Annals of Mathematics)

Abstract

This paper presents some considerations about the Goldbach's conjecture (GC). The work is based on elementary results of the number theory and it provides a constructive method that permits, given an even integer, to find at least a pair of prime numbers according to the GC. Eventually a correlation will be provided between this constructive method and a class of problems of operations research.

Keywords. Goldbach's Conjecture, Congruencies, Chinese remainder theorem, Operations research, Convex feasibility problem, Knapsack problem

1 Introduction

In his famous letter to Leonhard Euler dated 7 June 1742[14], Christian Goldbach stated his famous conjecture that "every integer greater than 2 can be written as the sum of three primes", statement that is equivalent to that every even number is a sum of two primes, as Euler stated in the letter dated 30 June 1742 ("I regard this as a completely certain theorem, although I cannot prove it"). This strong Goldbach conjecture implies the conjecture that all odd numbers greater than 7 are the sum of three odd primes, which is known today variously as the "weak" Goldbach conjecture, the "odd" Goldbach conjecture, or the "ternary" Goldbach conjecture.

In 1923 the ternary conjecture has been proved under the assumption of the truth of the generalized Riemann hypothesis[10], in 1937 Vinogradov[4] removed the dependence on the Riemann Hypothesis, and proved that this it true for all sufficiently large odd integers. In 1956 Borodzkin showed that an integer greater than a large integer is sufficient in Vinogradov's proof. In 1989 and 1996 Chen and Wang reduced this bound[7]. Using Vinogradov's method, Chudakov,van der Corput, and Estermann[5, 6, 9] showed that almost all even numbers can be written as the sum of two primes (in the sense that the fraction of even numbers which can be so written tends towards 1). In a very recent paper mathematician Terence Tao of the University of California, Los Angeles, has inched toward a proof, in fact he has shown that one can write odd numbers as sums of, at most, five primes [23].

Although believed to be true, the binary Goldbach conjecture is still lacking a proof. We state it as follows.

Statement 1.1. Every even integer greater than 4 can be written as the sum of two primes.

2 Alternative statement of the problem

Consider a natural number e, we can state the conjecture in term of the even natural number $2e$ as follows.

Statement 2.1. For every integer $e \geq 3$ there exists a couple of odd prime numbers q_1 and q_2 such that e is their semi-sum, that is: $e = \frac{q_1+q_2}{2}$.$\square$

This statement is equivalent to the original one considering that $2e$, with e a generic integer, is always an even natural, and that the original statement is verified for number 4 as it is the sum of $2 + 2$, with 2 the even prime.

Definition 2.1. Two prime numbers q_1 and q_2, with $q_2 > e > q_1$, are called *mirror primes respect to an integer* e if there exists an integer d such that: $e - q_1 = q_2 - e = d \geq 1$.

So another equivalent statement to 2.1, and of course to 1.1, is (see the figure):

Statement 2.2. For every natural $e \geq 3$ there exists a pair of mirror primes respect to e.$\square$



The statement 2.2, leads us to analyze when, given a natural number e, there exists at least one integer d defining two mirror primes respect to e. First of all we have to ask if and when there exist such number $d \geq 1$ (or, if it is taken as negative quantity, it must hold: $d \leq -1$). Moreover we are interested in an integer such that $e \pm d$ are primes (recall that the primality can be checked by means of an efficient algorithm currently available [1]).

3 Notation

Let observe that in the following we use that notation:

With the symbol $\pi(x)$ we denote the number of primes less than or equal to x.

With the expression $(n, m]$ we indicate the interval open on left and close on right.

With the expression $a \nmid b$ we denote that a doesn't divide b.

In some passages the symbol $[x]$ indicates the greatest integer less then or equal to x, but in general the square brackets are used to group symbols in an expression.

4 The Chinese Remainder Theorem and the number d

About the existence of prime numbers in the intervals definened by the number e, that is $(0, e)$ and $(e, 2e)$, the following result holds.

Lemma 4.1. Given an arbitrary integer $e > 3$ there is at least one prime number in the intervals $(0, e)$ and $(e, 2e - 2)$.

Proof. In the first interval we have trivially at least the two primes 2 and 3. Regarding the second interval, actually it is the Bertrand's postulate, also called the Bertrand-Chebyshev theorem or Chebyshev's theorem: it states that if $e > 3$, there is always at least one prime p between e and $2e - 2$ [15].□

Furthemore we can ask if there are particular relations among q_1, q_2, e, and d. We can easily observe that the mirror primes can not be prime factors neither of the natural number $2e$ nor of d. We summarize those remarks in the following Lemma.

Lemma 4.2. Given e, even integer number, for all pairs of mirror primes respect to e it holds:

(a) $q_1 \nmid e$;

(b) $q_2 \nmid 2e$;

(c) $q_1 \nmid d$.

Proof. We prove (a) by reductio ad absurdum. We have: $q_2 = 2e - q_1$, and if $q_1 | e$ then $q_1 | q_2$, but $q_2$is prime and so it is impossible. The statement (b) follows from the identity $q_1 = 2e - q_2$and from the fact that $q_1$is by hypothesis prime. Similarly the third statement follows from observing that $q_2 = q_1 + 2d$.□

Remembering that a prime number is a natural number greater than 1 that has no positive divisors other than 1 and itself, it is easily to infer that every composit number n has a prime factor less than or equal to $\sqrt{n}$. In fact if n has a prime factor less than or equal to its square root it is a composite number. Conversely let suppose for simplicity that n could be written, according to the fundamental theorem of arithmetic [14, 13], as the product of two primes a and b. If both of these two numbers were greater than $\sqrt{n}$ we should have that:

$n = a \cdot b > \sqrt{n} \cdot \sqrt{n} = n$, that is $n > n$, and it is not possible. The underlying rationale can be generalized to an arbitrary number of prime factors.

So at least one prime divisors of each of the two quantity $e \pm d$, if composite, are in the interval $\left[1, \sqrt{e \pm d}\right]$. On the other hand if $e \pm d$ are not divisible by any prime less then or equal to $\sqrt{e \pm d} < \sqrt{2e}$ then they can not be a composite integer, so it has to be prime.

The following lemma introduces the calculation method for the number d depending on an arbitrary choice of suitable integers b_i.

Lemma 4.3. Given a natural number e and a set of $k = \pi(\sqrt{2e})$ arbitrary integers b_i, there exists an integer number d solution of the following system of congruencies:

$$d \equiv b_i \ (mod\, p_i), \ i = 1..k$$

where p_i are the $k = \pi(\sqrt{2e})$ primes less then or equal to $\sqrt{2e}$.

Proof. Observing that $(p_i, p_j) = 1$ for $i \neq j$, the Chinese Remainder Theorem (CRT) [12] tell us that the system has a solution congruent to $p_1 \cdot p_2 \cdot ... \cdot p_k$.

The solution can be calculated by the following formula:

$d = \sum_{1 \leq i \leq k} b_i P_i P_i'$, where: $P = p_1 \cdot p_2 \cdot ... \cdot p_k$,$P_i = P/p_i$, and P_i' is the inverse of P_i, that is: $P_i P'_i \equiv 1(mod\, p_i)$.□

Let note that the right terms of several congruencies could be considered as negative integers, in this case the result of the lemma is still valid (that is, supposing $b_i > 0$, we would have: $d \equiv -1 \cdot b_i \ (mod\, p_i)$ for some $i \in [1..k]$, and the correspondent terms of the expession of the solution would be modified accordingly multiplying by -1).

Supposing that we can choose a set of integer numbers b_i with $i = 1..\pi\left(\sqrt{2e}\right)$ such that $b_i not \equiv \pm e \, (mod\, p_i)$, then the number d calculated by the foregoing method, if it is in the interval $[-(e-2), +(e-2)]$, is such that:

$d \equiv b_i \ (mod\, p_i), \, and \, b_i not \equiv \pm e \, (mod\, p_i) \, , \Rightarrow \, d\, not \equiv \pm e \, (mod\, p_i)$, that guarantees $e \pm d$ being a pair of primes.

5 How to calculate the integers b_i

The system of congruencies defined in the Lemma 3 provides an integer d that depends on the choice of the k integers b_i. Let us try to state the requirements they must meet. First of all we want the followings k conditions hold: $b_i not \equiv \pm e (mod\, p_i)$, in order to assure that $e \pm d \, not \equiv 0 \, (mod\, p_i)$, that is $p_i \nmid e \pm d$, which ensures that $e \pm d$ will be prime numbers. A possible choice for the number b_i could be that provided by the following lemma.

Lemma 5.1. Let e be a positive integer, we denote $p_1, p_2, ...p_k$ the first k primes with $k = \pi(\sqrt{2e})$, then the quantity defined as follows are not congruent to $\pm e$ moduli p_i for each i:

$b_i = \left[\frac{e}{p_i}\right] p_i$, if $p_i \nmid e$, and $b_i = \left[\frac{e}{p_i^{\alpha_i}}\right] p_i + 1$, if $p_i \mid e$ with $\alpha_i$the $i-th$ prime power according to the fundamental theorem of arithmetic[14, 13], and $[x]$ indicates the greatest integer less then or equal to x.

Proof. First consider the case in which $p_i \nmid e$. Observe that the expression $e - \frac{b_i}{p_i}$ represents the remainder after the division $\frac{e}{p_i}$, so it is not divisible by p_i. If $e \pm b_i$ was divisible by p_i we could write: $e = (h \mp \left[\frac{e}{p_i}\right])p_i$, with h a generic integer, but this would mean that $p_i \mid e$, contra hypothesis. Now suppose $p_i \mid e$, similarly if $e \pm b_i$ was divisible by p_i we could write: $e = (h \mp \left[\frac{e}{p_i^{\alpha_i}}\right])p_i + 1$, with h a generic integer, but but this would mean that $p_i \nmid e$, contra hypothesis.$\square$

Unfortunately the choice of the numbers b_i as indicated by lemma 5.1 doesn't assure that d is less than our integer e, let see for instance the following sidebar related to the integer 68.

$e = 68$

p_i	P_i	P'_i	b_i
2	1155	1	35
3	770	2	66
5	462	3	65
7	330	1	63
11	210	1	66

$d = 266805\,(mod\,2310) = 1155$. $p_1 = d - e = 1087$, $p_2 = d + e = 1223$, both prime numbers.

In order to avoid values of d greater than or equal to $e-2$ as well as, because the solution d could be negative depending on the choice of the sign of b_i, in order to avoid values less than or equal to $-(e-2)$, the following constraints must hold:

$$Q_k \cdot B^T \le e - 2$$

$$Q_k \cdot B^T \ge -(e - 2)$$

where: $B_{1\times k} = (\pm b_1 \;\pm b_2 \;\pm b_3 ... \pm b_k)$,

$$Q_{k\times k} = \begin{bmatrix} P_1 \cdot P'_1 & 0 & 0 & ... & 0 \\ P_1 \cdot P'_1 & P_2 \cdot P'_2 & 0 & ... & 0 \\ ... & ... & ... & ... & ... \\ P_1 \cdot P'_1 & P_2 \cdot P'_2 & P_3 \cdot P'_3 & ... & P_k \cdot P'_k \end{bmatrix}, \text{ with } det(Q) \neq 0.$$

Observe that the solution d of the system of congruencies of lemma 3 is $Q_k \cdot B^T$, where Q_k is the $k-th$ row of the matrix Q, where $k = \pi(\sqrt{2e})$.

Let define now a function $\kappa : (N^{+} \times N_{prime}) \to N$ as follows (where $N^{+} = N \setminus \{0\}$,$N_{prime} = \{n \geq 2, n\, prime\}$):

$$\kappa(x,p) = \begin{cases} 1, & p \nmid x \\ 2p, & p \mid x \end{cases}$$

Lemma 5.2. The definition of $\kappa(\cdot,\cdot)$ allow us to calculate at least one set of integer b_i such that $b_i not\equiv \pm e\ (mod\, p_i)$ with the following formula:

$$b_i = \min_{1 \leq j \leq 2p_i - 1} (\kappa(e+j,\, p_i) \cdot \kappa(e-j,\, p_i) \cdot j)$$

Proof. The expression provided assures that every b_i is not a multiple of $\pm e$ moduli the $i - th$ prime p_i. In fact the $j - th$ term of the set under analysis (let refer to it as b_i^j) is equal to j when p_i doesn't divide $e \pm j$ and it is equal to $2 \cdot j \cdot p_i$ or $4 \cdot j \cdot p_i^2$ otherwise. This leads to a partition in two subsets such that all the elements in the first set are less than all the elements in the second one, in particular we have:

$$\left\{b_i^j\right\} = \{b_i^m\} \cup \{b_i^n\}\,,\ m,n < 2p_i, \mid \{b_i^m\} \mid + \mid \{b_i^n\} \mid = 2p_i - 1, \{b_i^m\} \cap \{b_i^n\} = 0$$

$$b_i^m < b_i^n, \forall m,n$$

$$e \pm m\ not \equiv 0\ (mod\, p_i), \forall m$$

$$e \pm n \equiv 0\ (mod\, p_i), \forall n$$

The last two relations derive from the definition of the function κ, similarly the first relation is consequence of the fact that p_i doesn't divide $e \pm m$ so the number b_i^m is less than $2p_i$ for each m, conversely because p_i divides $e \pm n$ the number b_i^n is greater than or equal to $2p_i$ for each n. The existence of a minimum, which is in the set $\{b_i^m\}$, is ensured by the fact that the set is not empty due to the presence of at least one value of the index j not congruent to $\pm e$ moduli p_i when j takes values between 1 and $2p_i - 1$. So it holds:

$$e \pm b_i^m\ not \equiv 0\ (mod\, p_i), \forall m$$

$$e \pm b_i^n \equiv 0\ (mod\, p_i), \forall n$$

and therefore choosing the minimum in the first set it assures that they are not congruent to $\pm e$ moduli p_i for each i.□

After obtaining the candidates b_i for each i, we have to resolve the system of congruencies of lemma 3 in order to calculate d. According to the explicit expression of the solution, we must consider 2^k formulas, with $k = \pi(\sqrt{2e})$, in order to obtain a number d less than $e - 1$.

THEOREM 5.1. Given an arbitrary integer $e \geq 3$, there exists a choice of the quantities $\pm b_i$,with b_i calculated according to lemma 5.2, that satisfies the following relations:

$$Q_k \cdot B^T \leq e - 2$$

$$Q_k \cdot B^T \geq -(e-2)$$

where $d = Q_k \cdot B^T$, is the solution of the system of k congruencies:

$$d \equiv b_i \; (mod \, p_i)$$

such that the two integers $e - d$ and $e + d$ are prime numbers, that is there exists a pair of *mirror primes respect to the integer e*.

Proof. In order to calculate the quantity d according to the lemma 3, we have to calculate the set of b_i (positives by construction) as stated in lemma 5.2. Once obtained these positive integers, we have to attribute the appropriate sign to each one. The choice of the integer $+b_i$ versus $-b_i$ can be viewed as the choice of two sets of binary variable x_i and y_i, where:

$$x_i = \begin{cases} 0 & \Leftrightarrow y_i = 1 \Leftrightarrow we\,choose\; -b_i \\ 1 & \Leftrightarrow y_i = 0 \Leftrightarrow we\,choose\; +b_i \end{cases}$$

The constrains can be written as:

$$\sum_{1 \leq i \leq k} w_i \cdot x_i + \sum_{1 \leq i \leq k} (-w_i) \cdot y_i \leq e - 2$$

$$\sum_{1 \leq i \leq k} w_i \cdot x_i + \sum_{1 \leq i \leq k} (-w_i) \cdot y_i \geq -(e - 2)$$

where:

$$w_i = b_i \cdot P_i \cdot P_i', \; i = 1..k.$$

It may be observed that $y_i = 1 - x_i$, and then the preceding constrains, after substituting $y_i = 1 - x_i$, can be written as (in the following the symbol [.] represents just parentheses):

$$\sum_{1 \leq i \leq k} w_i \cdot x_i \leq \frac{1}{2} \left[(e-2) + \sum_{1 \leq i \leq k} w_i \right]$$

$$\sum_{1 \leq i \leq k} w_i \cdot x_i \geq \frac{1}{2} \left[-(e-2) + \sum_{1 \leq i \leq k} w_i \right]$$

Therefore it must be proved that, under the assumption that $e \geq 3$, the feasible set defined by the constraints is not empty. In other words, at least one choice of the binary variables x_i satisfies:

$$\sum_{1\le i\le k} w_i \cdot x_i \le \frac{1}{2}\left[(e-2)+\sum_{1\le i\le k} w_i\right] = \frac{1}{2}\sum_{1\le i\le k} w_i + E$$

$$\sum_{1\le i\le k} w_i \cdot x_i \ge \frac{1}{2}\left[-(e-2)+\sum_{1\le i\le k} w_i\right] = \frac{1}{2}\sum_{1\le i\le k} w_i - E$$

where $E = \frac{(e-2)}{2}$.

Now observe that the two constraints define a convex set. In fact, for all $\alpha \in (0,1)$ and considering two elements of this set x_i^1 and x_i^2 satisfying the constraints, we have:

$$\sum_{1\le i\le k} w_i \cdot (\alpha x_i^1 + (1-\alpha) x_i^2) = \alpha \sum_{1\le i\le k} w_i \cdot x_i^1 + (1-\alpha)\sum_{1\le i\le k} w_i \cdot x_i^2 \le \frac{1}{2}\sum_{1\le i\le k} w_i + E$$

$$\sum_{1\le i\le k} w_i \cdot (\alpha x_i^1 + (1-\alpha) x_i^2) = \alpha \sum_{1\le i\le k} w_i \cdot x_i^1 + (1-\alpha)\sum_{1\le i\le k} w_i \cdot x_i^2 \ge \frac{1}{2}\sum_{1\le i\le k} w_i - E$$

The problem to find a solution in this set is known in literature as "Convex Feasibility Problem" [18, 19, 20]. It consists in finding a point in the intersection of convex sets. The common way for solving it is the relaxation-projection algorithm [21, 22]. Howevever we are interested for our goal in proving the existence of a choice of the variables x_i. Let observe that we may order the terms w_i such that: $w_i \le w_{i+1}, \forall i = 1..k-1$, and then we can determine the index h such that:

$$\sum_{1\le i\le h} w_i < \frac{1}{2}\sum_{1\le i\le k} w_i$$

$$\sum_{1\le i\le h+1} w_i \ge \frac{1}{2}\sum_{1\le i\le k} w_i$$

and therefore choose the variables as follows:

$$x_i = 1,\ i = 1..h,\ x_i = 0,\ i = h+1..k$$

With this choice of the values of the k variables we may have (note that $E > 0$ and $k \ge 2$ if $e \ge 3$):

$$\frac{1}{2}\sum_{1\le i\le k} w_i > \sum_{1\le i\le h} w_i = \sum_{1\le i\le k} w_i \cdot x_i > \sum_{1\le i\le k} w_i \cdot x_i - E$$

And therefore the first constraint is satisfied.

Regarding the second constraint let notice that from the expression provided by lemma 5.2 and observing that $p_i \le \sqrt{2e}$, the following holds:

$$b_i \leq 2p_i - 1 \leq 2\sqrt{2e} - 1$$

Furthemore from the fact that $P'_i \leq p_i$, considering the expression provided by lemma 4.3, it holds:

$$P_i P'_i \leq \pi(\sqrt{2e})\sqrt{2e}$$

From these two inequalities it is now possible express a general upper bound for the terms w_i:

$$w_i \leq (2\sqrt{2e} - 1)\pi(\sqrt{2e})\sqrt{2e} \leq 4e\pi(\sqrt{2e})$$

We can write indeed:

$$\frac{1}{2}\sum_{1\leq i\leq k} w_i - \sum_{1\leq i\leq k} w_i \cdot x_i \leq$$

$$\leq 2ek\pi(\sqrt{2e}) - 4eh\pi(\sqrt{2e}) = 2e\pi(\sqrt{2e})(k - 2h)$$

Conseguently the second inequality under analysis can be written as:

$$2e\pi(\sqrt{2e})(k - 2h) \leq \frac{e-2}{2}$$

that is:

$$2 + 4e\pi(\sqrt{2e})(k - 2h) \leq e$$

expression satisfied if:

$$h \geq \frac{k}{2}$$

Condition that is necessary true based on the definition of index h.

Then the particular choice of the set of binary variables satisfies both of the constraints.□

Let see as an example the following sidebar where the calculus is provided for the integer 16.

$$e = 16$$

p_i	j	b_i^j	min
2	$[1, 3]$	$\{1, 32, 3\}$	1
3	$[1, 5]$	$\{6, 12, 3, 24, 30\}$	3
5	$[1, 9]$	$\{10, 2, 3, 40, 5, 60, 7, 8, 90\}$	2

from the CRT:

p_i	P_i	P'_i	b_i
2	15	1	1
3	10	1	3
5	6	1	2

$$d_h = \pm b_1 \cdot 15 \pm b_2 \cdot 10 \pm b_3 \cdot 6, 1 \leq h \leq 2^3$$

for example consider the following choices:

$$+1 \cdot 15 + 3 \cdot 10 + 2 \cdot 6 = 57 \geq e - 2$$

$$-1 \cdot 15 + 3 \cdot 10 - 2 \cdot 6 = 3 \leq e - 2$$

$$+1 \cdot 15 - 3 \cdot 10 + 2 \cdot 6 = -3 \geq -(e - 2)$$

$$-1 \cdot 15 - 3 \cdot 10 + 2 \cdot 6 = -33 < -(e - 2)$$

... ...

the second and the third are feasible solutions

therefore if we can choose $d = 3$:

$$16 - 3 = 13 \text{ , } prime$$

$$16 + 3 = 19 \text{ , } prime$$

so that:

$$13 + 19 = 32 = 2 \cdot 16$$

While we have spoken up to now in terms of deriving a generic value for the quantity d, it could be required to identify the greatest or the smallest d with the properties we have dicussed in the preceding. The natural way to do this is to write the related optimization problem in terms of **integer linear program**, and we can see that it is a special case of well known formulation classified in literature as 'Knapsack Problem'[16]:

maximize (or minimize) $W_{(1\times k)}X_{(k\times 1)}$
subject to:
$WX \leq U$
$-WX \leq U'$
$x_i \in \{0, 1\}$

where:
$U = \frac{1}{2}\sum\limits_{1\leq i\leq k} w_i + E$, $U' = -\frac{1}{2}\sum\limits_{1\leq i\leq k} w_i + E$,
$W_{(1\times k)} = (w_1 \ldots w_k)$, $X_{(k\times 1)} = (x_1 \ldots x_k)^T$.

Given a set of items, each with a weight and a value (in our case they are coincident and the problem is called 'Subset sum problem'), determine the number of each item to include in a collection so that the total weight is less than or equal to a given limit and the total value is as large as possible. It derives its name from the problem faced by someone who is constrained by a fixed-size knapsack and must fill it with the most valuable items. Moreover, considering the two sets of constrains, we observe that various methods are known in literature in order to deal with negative weight [17].

References

[1] Manindra Agrawal, Neeraj Kayal, Nitin Saxena, *Primes is in P*, Volume 160, Issue 2 (2004), Pages 781-793.

[2] Christian Goldbach, *Lettre XLIII*, `http://www.math.dartmouth.edu/~euler/correspondence/letters/OO0765.pdf`.

[3] Eric W.Weisstein, *Goldbach Conjecture*, `http://mathworld.wolfram.com/`.

[4] I.M. Vinogradov. *The Method of Trigonometrical Sums in the Theory of Numbers.* New York: Interscience (1954).

[5] Nikolai G. Chudakov, *On the Goldbach problem*, Doklady Akademii Nauk SSSR 17 (1937): 335–338.

[6] J. G. Van der Corput, *Sur l'hypothèse de Goldbach*, Proc. Akad. Wet. Amsterdam 41 (1938): 76–80.

[7] J. R. Chen and Y. Wang, *On the odd Goldbach problem*, Acta Math. Sinica, 32 (1989), 702–718.

[8] Chen, Jing Run and Wang, Tian Ze, *The Goldbach problem for odd numbers*, Acta Math. Sinica (Chin. Ser.), 39:2 (1996) 169–174.

[9] T. Estermann, *On Goldbach's problem: proof that almost all even positive integers are sums of two primes*, Proc. London Math. Soc.. 2 44: 307–314 (1938).

[10] G. H. Hardy and J. E. Littlewood, *Some problems of 'partitio numerorum' : III: on the expression of a number as a sum of primes*, Acta Math., 44, 1-70 (1923). Reprinted in "Collected Papers of G. H. Hardy," Vol. I, pp. 561-630, Clarendon Press, Oxford, 1966.

[11] J.R Chen, *On the representation of a large even integer as the sum of a prime and the product of at most two primes*, J. Kexue Tongbao 17 (1966), 385–386.

[12] Tom M. Apostol, *Introduction to Analytic Number Theory*, (1976) Springer-Verlag, New York.

[13] Euclid's Element, Book VII, propositions 30 and 32.

[14] Gauss, Carl Friedrich; Clarke, Arthur A. (translator into English) (1986), *Disquisitiones Arithemeticae* (Second, corrected edition), New York: Springer.

[15] P. Chebyshev *Mémoire sur les nombres premiers*, Mém. Acad. Sci. St. Pétersbourg 7, 17-33, (1850) 1854. Reprinted as §1-7 in Œuvres de P. L. Tschebychef, Tome I. St. Pétersbourg, Russia: Commissionaires de l'Academie Impériale des Sciences, pp. 51-64, 1899.

[16] Michael R. Garey, David S. Johnson, *Computers and Intractability: A Guide to the Theory of NP-Completeness.* W.H. Freeman. (1979).

[17] F. Glover, *A multiphase-Dual Algorithm for the Zero-One Integer Programming Problem*, Operations Research, Vol.13, N.16 (Nov-Dec, 1965).

[18] F. E. Browder, "Convergence theorems for sequences of nonlinear operators in Banach spaces," Mathematische Zeitschrift, vol. 100, no. 3, pp. 201-225, July 1967.

[19] F. E. Browder, "Convergence theorems for sequences of nonlinear operators in Banach spaces," Mathematische Zeitschrift, vol. 100, no. 3, pp. 201-225, July 1967.

[20] Maruster, St., Popirlan, C., On the regularity condition in a convex feasibility problem, Nonlinear Analysis, Vol. 70 (2009), pp. 1923- 1928.

[21] Agmon, S., The relaxation method for linear inequalities, Canad. J. Math., Vol. 6 (1954), pp. 382-392.

[22] [3] Motzkin, T.S., Schoenberg, I.J.,"The relaxation method for linear inequalities", Canad. J. Math., Vol. 6 (1954), pp. 393-404.

[23] T.Tao, "Every odd number greater than 1 is the sum of at most five primes", eprint arXiv:1201.6656, 01/2012.